



Igor Grosman, DO; F.A.S.G
 Alexander Brun, MD
 Prateek Chapalamadugu, MD
 Nancy Chen, MD
 Konstantin Peysin, DO
 Yevgeniy Mikityanskiy, DO

Board Certified in Gastroenterology & Hepatology
 E – Mail: info@TriboroughGI.com, www.TriboroughGI.com

Main Office 1672-1674 Sheepshead Bay Rd Brooklyn, NY 11235 T- (718) 332 - 0600	Endoscopy Center 214 Avenue P (West 6 th & 7 th St) Brooklyn, NY 11204 T- (718) 339 - 5678	Mount Sinai Hospital Brooklyn 3201 Kings Highway Brooklyn, NY 11234 T- (718) 951 – 2801	New York Community Hospital 2525 Kings Highway Brooklyn, NY 11229 T- (718) 692-5343	Staten Island Office 2076 Hylan Blvd, Suite#2 Staten Island, NY 10306 T- (718) 987-1700	Statcare 2063 Bartow Ave Bronx, NY 10475 T- (718) 332 - 0600
---	---	---	---	---	--

Colonoscopy – Golytely

APPOINTMENT INFORMATION

Patient Name: _____

Date: _____

Time: _____

Surgical Center Appointment Cancellation Policy

We **require** that you give our office **24 Hours'** notice in the event that you need to **reschedule your appointment**. This allows for other patients to be scheduled into that appointment. If you miss an appointment without contacting our office within the required time, this is considered a missed appointment. **A FEE OF \$100 WILL BE CHARGED TO YOU.**

GOLYTELY PREP INSTRUCTIONS:

1 WEEK PRIOR

- Fill your prescription for Golytely and 2 tablets of Bisacodyl (Dulcolax) at the pharmacy.
- Arrange for a driver to take you home on the day of the procedure.
- If you have diabetes, check with your doctor regarding diet and medication instructions.
- Please **DISCONTINUE** fiber supplements and iron supplements including multivitamins and blood thinners (**Coumadin and Plavix**).
- **DO NOT STOP ASPIRIN** unless doctor told you to do so.

1 DAY PRIOR

- Start a clear liquid diet in the morning. **NO SOLID FOOD (NO BREAKFAST, LUNCH, or DINNER).**
Clear liquids include: Broths, Water, Clear fruit juices (apple or white), Soda, Gatorade, Jell-O, Popsicles, Coffee or Tea (No Milk or Cream). You may use sugar or lemon. **Please do not have any dairy products, or Orange Juice or Anything Red**
- Take 2 tablets of Bisacodyl at 5PM.
- Mix Golytely solution as directed on the container. You may put it in the refrigerator to chill before drinking.
- Between 5PM and 8PM start drinking one 8oz glass of the solution every 15 minutes.
- Nothing is to be taken after midnight prior to the procedure.

PROCEDURE DAY

- You may take your morning medications (**High Blood Pressure**) with a small sip of water.
- If you have asthma, please bring your inhaler with you.

A RESPONSIBLE ADULT MUST ACCOMPANY ALL PATIENTS THE DAY OF THE PROCEDURE. NO PATIENT SHOULD DRIVE ON THE DAY OF THE COLONOSCOPY.

*****NOTE: ALL PATIENT'S MUST DO COVID-19 SWAB TEST 5 DAYS BEFORE PROCEDURE DATE*****